# Biotype-dependent secondary symbiont communities in sympatric populations of *Bemisia tabaci*

# E. Chiel<sup>1,2</sup>, Y. Gottlieb<sup>3</sup>, E. Zchori-Fein<sup>1\*</sup>, N. Mozes-Daube<sup>1</sup>, N. Katzir<sup>4</sup>, M. Inbar<sup>2</sup> and M. Ghanim<sup>3</sup>

<sup>1</sup>Department of Entomology, Newe-Ya'ar Research Center, ARO, Ramat-Yishai, 30095: <sup>2</sup>Department of Evolutionary and Environmental Biology, University of Haifa, Haifa, 31905: <sup>3</sup>Department of Entomology, the Volcani Center, ARO, Beit-Dagan: <sup>4</sup>Department of Vegetable Crops, Newe-Ya'ar Research Center, ARO, Ramat-Yishai, 30095

# Abstract

The sweet potato whitefly, Bemisia tabaci, harbors Portiera aleyrodidarum, an obligatory symbiotic bacterium, as well as several secondary symbionts including Rickettsia, Hamiltonella, Wolbachia, Arsenophonus, Cardinium and Fritschea, the function of which is unknown. Bemisia tabaci is a species complex composed of numerous biotypes, which may differ from each other both genetically and biologically. Only the B and Q biotypes have been reported from Israel. Secondary symbiont infection frequencies of Israeli laboratory and field populations of B. tabaci from various host plants were determined by PCR, in order to test for correlation between bacterial composition to biotype and host plant. Hamiltonella was detected only in populations of the B biotype, while Wolbachia and Arsenophonus were found only in the Q biotype (33% and 87% infection, respectively). Rickettsia was abundant in both biotypes. Cardinium and Fritschea were not found in any of the populations. No differences in secondary symbionts were found among host plants within the B biotype; but within the Q biotype, all whiteflies collected from sage harboured both Rickettsia and Arsenophonus, an infection frequency which was significantly higher than those found in association with all other host plants. The association found between whitefly biotypes and secondary symbionts suggests a possible contribution of these bacteria to host characteristics such as insecticide resistance, host range, virus transmission and speciation.

Keywords: Arsenophonus, Hamiltonella, host plant, Rickettsia, whitefly, Wolbachia

#### Introduction

The sweet potato whitefly, *Bemisia tabaci* (Gennadius), is an exceedingly polyphagous pest capable of developing on hundreds of plant species, including many agricultural crops (Oliveira *et al.*, 2001). Its broad host range has been the subject of many studies aimed at understanding the

\*Corresponding author E-mail: einat@agri.gov.il whitefly's host selection preferences, as well as the influence of host plants on whitefly fitness (Van Lenteren & Noldus, 1990; Gerling & Mayer, 1996).

The taxonomic status of *B. tabaci* is a complex issue; different populations, while morphologically indistinguishable, display a high degree of genetic and biological variability, and this has led to the definition of *B. tabaci* as a species complex. To date, more than 20 biotypes, differing genetically and in host range, insecticide resistance, and ability to transmit plant viruses and cause plant disorders, have been designated (Brown *et al.*, 1995; Perring, 2001).

While some of the biotypes have a limited host range and local distribution, others, particularly biotype B, are widely distributed and polyphagous. Because biotypes can not be distinguished by morphological characters, their identification is dependent on molecular techniques.

De Barro et al. (2005) recently proposed grouping B. tabaci populations into six geographical races based on COI and ITS1 gene sequences, but the common nomenclature used is still that of biotypes. In Israel, two biotypes have been reported: B (also referred to as the silverleaf whitefly, B. argentifolii Bellows & Perring); and Q. According to the classification of De Barro et al. (2005), these two biotypes belong to two closely related sister groups: the Q biotype, which originated in the Mediterranean Basin; and the B biotype, which arose in the Middle East and Asia Minor. Despite their close geographical origins, these two biotypes rarely interbreed. Females have been reported to reject the mating attempts made by males, and the rare interbiotype crosses observed resulted in either sterile F1 or a hybrid breakdown at the F2 (Ma et al., 2004; Shai Morin, personal communication). Both pre- and post-zygotic reproductive barriers, thus, seem to exist between the Q and B biotypes tested. The most prominent phenotypic differences between the two biotypes are the higher fecundity and competitive abilities of the B biotype and the greater degree of pesticide resistance of the Q biotype (Pascual & Callejas, 2004; Horowitz et al., 2005).

Like other phloem-feeders, B. tabaci harbors primary symbionts, obligatory bacteria which are essential for its survival and development. The symbiont, Portiera aleyrodidarum, is confined to specialized cells (bactriocytes) and is vertically transmitted (reviewed in Baumann, 2005). In addition to this primary symbiont, B. tabaci is host to a diverse array of secondary symbionts (SS). Although SS are facultatively associated with their hosts, i.e. they are not essential for host survival, they may play important roles in their host's ecology and evolution. SS currently described from B. tabaci are Arsenophonus, Hamiltonella, Wolbachia, Cardinium, Fritschea and Rickettsia (reviewed in Baumann, 2005; Gottlieb et al., 2006), but only Hamiltonella and Rickettsia have been detected in Israeli B. tabaci populations (Gottlieb et al., 2006). The function of these SS in the whitefly is unknown, but other insect-SS systems reveal assorted and intriguing phenomena. In the pea aphid, Acyrthosiphon pisum, for example, five different SS have been identified and studied. Two of them, Hamiltonella defensa and Serratia symbiotica, increase resistance to parasitoids of the genus Aphidius (Oliver et al., 2003; Ferrari et al., 2004). S. symbiotica also improves the aphid's heat tolerance (Montllor et al., 2002) and compensates for the absence of the primary symbiont (Koga et al., 2003). Another SS, Regiella insecticola, is strongly correlated with host plant range, susceptibility to environmental factors, resistance to entomophthogenic fungi and the alteration of dispersal and mating (Tsuchida et al., 2002, 2004; Scarborough et al., 2005; Leonardo & Mondor, 2006).

Bacteria of the genus, *Rickettsia* ( $\alpha$ -Proteobacteria), are mostly known as obligate intracellular symbionts of bloodfeeding arthropods and as causative agents of many vertebrate diseases such as typhus. They are evidently associated with various phenomena in non-blood feeding arthropods as well, including sex ratio disorders (reviewed in Perlman *et al.*, 2006). *Wolbachia* ( $\alpha$ -Proteobacteria) is a wellknown 'reproductive manipulator', owing to its ability to alter many arthropods' reproduction processes (Stouthamer et al., 1999). Wolbachia was traced in previous surveys of B. tabaci (Zchori-Fein & Brown, 2002; Nirgianaki et al., 2003; De Barro, 2005), and it has been cited as the possible cause of cytoplasmic incompatibility among some B. tabaci biotypes (De-Barro & Hart, 2000). Arsenophonus spp. (y-Proteobacteria) also are distributed among a variety of insects including whiteflies, aphids, psyllids and a louse fly (Baumann, 2005; Dale et al., 2006). A. nasoniae causes male embryo mortality in the parasitoid wasp Nasonia vitripennis (Gherna et al., 1991), but its effect on other hosts is unknown. Cardinium (Bacteroidetes) is also known as a reproductive manipulator (Weeks et al., 2003). Bacteria of the order, Chlamydiales, are often pathogenic to humans and vertebrates, and some are vectored by arthropods (http:// www.chlamydiae.com). However, nothing is known about the phenotype of Fritschea (Chlamydiales) in B. tabaci (Everett et al., 2005).

This study was initiated in order to determine if SS associations and infection frequencies are correlated with the biotype and host plant of *B. tabaci*, in both field and laboratory populations. We see this as a first step towards understanding the phenotypical characteristics of these symbionts in this insect species complex.

# Methods and materials

### Whitefly origin and rearing

Samples of *B. tabaci* were obtained from the laboratory collection of Dr R. Horowitz (Beit Dagan, Israel). These whiteflies were originally collected in various agricultural crops, during various years and seasons, at different locations in Israel (table 1) and were maintained in the laboratory as separate populations on cotton seedlings ('Acala') ( $26 \pm 2^{\circ}$ C, 60% RH, 14L:10D). In order to screen field populations, live *B. tabaci* adults were collected from various crops at 18 different locations throughout Israel (table 2) during September and October 2005 and were immediately preserved in 96% ethanol.

# Screening for the presence of secondary symbionts

Laboratory populations were first screened in bulk to determine the presence of the various SS. Twenty to fifty adult whiteflies from each lab population were ground together in a lysis buffer as described by Frohlich *et al.* (1999). This was followed by incubation at 65°C for 15 min, and then at 95°C for 10 min. The presence of *Rickettsia, Hamiltonella, Wolbachia, Cardinium, Arsenophonus* and *Fritschea* in the samples was determined using genusspecific primers for amplifying 16S or 23S rDNA gene fragments (table 3). Reactions were carried out in 25 µl volumes each containing 2µl of the template DNA lysate, 10 pM of each primer, 0.2 mM dNTP's, 1× RedTaq buffer and one unit of RedTaq DNA polymerase (Sigma). PCR products were visualized on 1.2% agarose gel containing ethidiumbromide.

In order to verify the products' identity, bands were eluted, cloned into the pGEM T-Easy plasmid vector (Promega) and transformed into *E. coli*. For each bacterium, two colonies were randomly picked and sequenced (ABI 3700 DNA analyzer, Macrogen Inc., Korea). The resulting

Laboratory population	Plant	Biotype	SS presence				
number			Н	А	W	R	
1	Abutilon	В	+	_	_	_	
2	Cotton	В	+	_	_	+	
3	Cotton	В	+	_	_	+	
4	Cotton	В	+	_	_	+	
5	Cotton	В	+	_	_	+	
6	Cotton	В	+	_	_	_	
7	Cotton	В	+	_	_	+	
8	Cotton	В	+	_	_	+	
9	Cotton	В	+	_	_	+	
10	Cotton	В	+	_	_	+	
11	Eggplant	В	+	_	_	+	
12	Eggplant	В	+	_	_	+	
13	Melon	В	+	_	_	+	
14	Pepper	В	+	_	_	+	
15	Potato	В	+	_	_	_	
16	Potato	В	+	_	_	+	
17	Watermelon	В	+	_	_	_	
18	Zucchini	В	+	_	_	_	
19	Basil	Q	_	+	_	+	
20	Cotton	Q	_	+	+	_	
21	Cotton	Q	_	+	+	+	
22	Cotton	Q	_	+	+	_	
23	Cotton	Q	_	+	_	+	
24	Cotton	Q	_	+	+	+	
25	Gerbera	Q	_	+	_	+	
26	Roses	Q	_	+	+	+	
27	Sunflower	Q	_	+	_	+	
28	Sunflower	Q	_	+	_	+	

Table 1. Presence (+) or absence (-) of secondary symbionts in *Bemisia tabaci* laboratory populations.

R, Rickettsia; H, Hamiltonella; W, Wolbachia; A, Arsenophonus.

Table 2. Infection frequencies of different secondary symbiont combinations in Bemisia tabaci field populations.

Field population	Host plant	n*	SS combination (%)										
number			B biotype		Q biotype								
			RH	Н	R	RWA	RW	RA	R	WA	W	А	None
1 2	Cotton	20 19	10	_	-	5	5	30 53	15 16		10	10	15
3		20 20	10	10	_	20 25	10 5	20 40	- 5	5 10		20 15	5
5 6	Cucumber	20 20	45 50	30 50	_	15	-	5	_	-	_	5	_
7	Eggplant	20	75	20	_	5	-	_	_	-	_	_	-
8 9	Lettuce	19 20	26 50	5 50	-	11 _		26	-	16	-	16	
10 11	Sage	20 20		-	_	30 35		70 55	-	-	-	-	
12	Squash	20	40	40	_	-	-	5	_	-	_	15	-
13 14 15 16	Sweet pepper	20 20 19 20	60 95 37 60	40 5 58 35	- - 5	- - -	- - -	- - 5 -	- - -	- - -	- - -	- - -	- - -
17 18	Tomato	20 18	75 11	25	_ _		-	33	-	- 6	_ 6	22	_ 6

n = total number of individuals tested from each population (including both biotypes).

R, Rickettsia; H, Hamiltonella; W, Wolbachia; A, Arsenophonus; -, zero.

Targeted gene	Primers' name	Primers sequence $(5' \rightarrow 3')$	Annealing temp. (°C)/ Product size (bp)	Reference
<i>Rickettsia</i> 16S rDNA	Rb-F Rb-R	GCTCAGAACGAACGCTATC GAAGGAAAGCATCTCTGC	60°/~900	Gottlieb et al., 2006
<i>Hamiltonella</i> 16S rDNA	Ham-F Ham-R	TGAGTAAAGTCTGGAATCTGG AGTTCAAGACCGCAACCTC	60°/~700	Zchori-Fein & Brown, 2002
Wolbachia 16S rDNA	Wol16S-F Wol16S-R	CGGGGGAAAAATTTATTGCT AGCTGTAATACAGAAAGTAAA	55°/~700	Heddi et al., 1999
Arsenophonus 23S rDNA	Ars23S-1 Ars23S-2	CGTTTGATGAATTCATAGTCAAA GGTCCTCCAGTTAGTGTTACCCAAC	60°/~600	Thao & Baumann, 2004
<i>Cardinium</i> 16S rDNA	CFB-F CFB-R	GCGGTGTAAAATGAGCGTG ACCTMTTCTTAACTCAAGCCT	58°/~400	Weeks et al., 2003
Fritschea 23S rDNA	U23F 23SIGR	GATGCCTTGGCATTGATAGGCGATGAAGGA TGGCTCATCATGCAAAAGGCA	60°/~600	Everett et al., 2005
B. tabaci MtCOI	C1-J-2195 L2-N-3014	TTGATTTTTTGGTCATCCAGAAGT TCCAATGCACTAATCTGCCATATTA	52°/~800	Khasdan et al., 2005

Table 3. PCR primers and conditions used in the study.

sequences were compared to known sequences in the databases using the BLAST algorithm in NCBI.

In order to assess within- and between-population infection rates of the bacteria found in the laboratory screening, 20 adults were taken from each field population and ground individually according to the method described above. They were screened with the relevant primers for the presence of the SS identified by the bulk screening of laboratory populations (table 3).

#### Identification of B. tabaci biotypes

Biotypes were identified using a method described by Khasdan *et al.* (2005). Briefly, in this method a fragment of the mtCOI gene is amplified by PCR (Frohlich *et al.*, 1999), followed by restriction endonuclease digestion with *VspI*, which generates a clear polymorphism between biotypes B and Q. PCR reactions were carried out as described above with the addition of 0.2 mM MgCl<sub>2</sub>. The PCR products were incubated with 0.1 units of *VspI* (Fermentas) at 37°C for 16 h before loading onto agarose gel.

#### Statistical analysis

SS frequencies among biotypes and/or plants were compared by a Pearson non-parametric test using the  $\chi^2$  distribution. JMP 5.0.1 software (SAS Institute) was used for all statistical analyses.

#### Results

#### Screening for the presence of secondary symbionts

# Laboratory populations

*Hamiltonella* was detected in all 18 B biotype populations, but was not found in any of the ten Q biotype populations. *Wolbachia* and *Arsenophonus* were found only in Q biotype populations (5/10 and 10/10, respectively). *Rickettsia* was found in both Q biotype (80%) and B biotype (72%) populations ( $\chi^2_1$ =0.21, *P*=0.65). Two Q populations were double-infected with *Arsenophonus* and *Wolbachia*, five were infected with both *Arsenophonus* and *Rickettsia*, and three



Fig. 1. Frequency of secondary symbiont combinations within field-collected individuals of B and Q biotypes of *Bemisia tabaci*. R, *Rickettsia*; H, *Hamiltonella*; W, *Wolbachia*; A, *Arsenophonus*;  $\blacksquare$ , B (n = 205);  $\square$ , Q (n = 150).

harbored all three SS (table 1). None of the populations screened tested positive for the presence of either *Cardinium* or *Fritschea*.

# Field populations

Since neither *Cardinium* nor *Fritschea* were detected in the mass laboratory screening, field populations were not screened for these two SS. As in the lab populations, *Hamiltonella* was detected in virtually all B biotype individuals, whereas none of the Q biotype whiteflies were infected with this bacterium (fig. 1). *Wolbachia* and *Arsenophonus* were detected only in Q biotype individuals (33% and 87% infection, respectively). Both biotypes harbored *Rickettsia*, with a somewhat higher infection percentage in the Q biotype (74% and 64%, respectively,  $\chi^2_1$ = 3.7, *P*=0.054). All possible combinations of *Rickettsia*, *Wolbachia* and *Arsenophonus* were found among the Q biotype individuals

(fig. 1); five of these individuals (3.3%) did not carry any of the SS, while none of the B individuals was SS-free ( $\chi^2_1$ = 6.93, *P* < 0.01). The infection categories distribution was significantly different across the two biotypes ( $\chi^2_9$ = 351.4, *P* < 0.0001).

The B biotype was dominant in field populations sampled from sweet pepper, cucumber, squash and eggplant (75–100%), while the Q biotype dominated in cotton and sage plots (80–100%) ( $\chi^2_7$  = 208.6, *P* < 0.0001). In tomato and lettuce, the results were not uniform (table 2).

Infection status of SS did not differ among host plants within the B biotype ( $\chi^2_{12}$ =8.6, *P*=0.73) but did vary within the Q biotype ( $\chi^2_{21}$ =42, *P*<0.01). The differences among host plants within the Q biotype were mostly associated with individuals collected from sage plots (*n*=38), all of which carried both *Rickettsia* and *Arsenophonus*. About one-third of these individuals also carried *Wolbachia*. Among individuals from other host plants, only ~38% carried both *Rickettsia* and *Arsenophonus* and 14–19% were triple infected (table 2). Infection with *Rickettsia* alone and with the combination of *Rickettsia* plus *Wolbachia* was found only in individuals sampled from cotton plots (seven and five individuals, respectively). *Wolbachia* infection rates were similar, 31–38% across samples from all host plants ( $\chi^2_3$ =0.26, *P*=0.96).

#### Discussion

#### Biotype and SS

Our study demonstrates a clear association between certain SS and *B. tabaci* biotypes. All Israeli populations of the B biotype harbor *Hamiltonella*, whereas *Wolbachia* and *Arsenophonus* are found exclusively in the Q biotype (fig. 1). Variability of symbiont combinations has been reported in the past for the A and B biotypes (Costa *et al.*, 1995), although at that time the identity of the bacteria was not established. It thus appears that biotype-dependent differences of SS composition can be of use in differentiating between *B. tabaci* biotypes.

Secondary symbionts are usually considered nonessential to their hosts, hence their presence between and within populations can be variable. Hamiltonella, for example, was previously reported from 40% of B. tabaci populations (Zchori-Fein & Brown, 2002) and from 0--46% of screened pea aphid populations. The incidence of Rickettsia in A. pisum also ranged from 1-48% in various reports (Chen et al., 2000; Darby et al., 2001; Tsuchida et al., 2002; Darby et al., 2003; Haynes et al., 2003; Ferrari et al., 2004). Exceptionally, all 40 clones of the aphid Uroleucon ambrosiae collected throughout the USA were found to carry Hamiltonella (Sandstrom et al., 2001), a result similar to the virtually 100% occurrence of Hamiltonella in the B biotype of B. tabaci reported here. Such a high incidence may indicate an obligatory or even mutualistic interaction between Hamiltonella and its host. There are many ways in which Hamiltonella may contribute to the whitefly. Two possibilities include complementing the primary symbiont, Portiera, as reported recently from a sharpshooter (Wu et al., 2006) and facilitating the higher fecundity of the B biotype, as compared to the Q and other biotypes (Brown et al., 1995; Nombela et al., 2001; Pascual & Callejas, 2004), thereby enhancing its own transmission. Hamiltonella might also be involved in plant physiological disorders that are exclusively caused by B biotype (e.g. squash silver-leafing). These hypotheses can be tested if *Hamiltonella*-free B biotype populations can be successfully established.

Wolbachia and Arsenophonus were harbored only by the Q biotype of *B. tabaci. Wolbachia* can be found in all major insect orders in varying frequencies (Stouthamer *et al.*, 1999; Werren & Windsor, 2000). The  $\sim$ 30% infection proportion of *Wolbachia* found in our study is in agreement with previous reports (Zchori-Fein & Brown, 2002; Nirgianaki *et al.*, 2003). Nirgianaki *et al.* (2003) reported that 11 out of 39 *B. tabaci* populations collected worldwide were infected with *Wolbachia*; of those, ten were 'non-silverleafing' biotypes, i.e. non-B biotype, as in our study. *Rickettsia* is the only SS that was found in both biotypes, a fact that might hint at a horizontal transmission pathway of this bacterium, possibly via the host plant.

The biotype-based divergence of SS resembles the strong association found between host races of *A. pisum* and their SS by Simon *et al.* (2003). In that study, *Hamiltonella* was found only in the alfalfa race of the aphid, while *Rickettsia* and *Serratia* were abundant in the pea race and rare on alfalfa and clover races. Double infections were also quite common in that study, as in ours. Other studies have revealed that *Rickettsia* and *Serratia* can suppress *Buchnera*, the primary symbiont of *A. pisum*, but *Serratia* can compensate for the essential role of *Buchnera* when the latter is absent (Koga *et al.*, 2003; Sakurai *et al.*, 2005). A quantitative PCR study can clarify if similar interactions exist in the *B. tabaci-SS* system.

There are several ways in which the biotype-dependence of SS communities can be explained:

**1.** Allopatric divergence. It has been suggested that historic geographic isolation has led to separation of races (De Barro *et al.*, 2005); accordingly, the B and Q biotypes acquired different sets of SS when they were isolated in the past, and only later became sympatric.

**2.** Sympatric divergence. (i) B and Q biotypes may have acquired different SS while in sympatry due to the genetic differences between them; or (ii) a SS (e.g. *Hamiltonella* or *Arsenophonus*) may have invaded one population, thus erecting a reproductive barrier between this population and others, resulting in the development of a new biotype.

Wolbachia, Rickettsia and Arsenophonus are known to cause reproductive manipulations in arthropods, but their role in B. tabaci is unknown. The possibility that these symbionts and others may cause reproductive isolation among biotypes via cytoplasmic incompatibility or other mechanisms should be carefully explored. Understanding the role of SS in the reproductive isolation of biotypes may help resolve the longstanding debate surrounding the taxonomic status of members of the B. tabaci complex; if the B and Q biotypes are reproductively isolated merely because of transient SS infections, they can be regarded as biotypes of the same species, while the lack of such influence will support their division into two distinct species. However, it should be noted that the ability of biotypes to interbreed may have been lost due to molecular distance, as proposed by De Barro et al. (2005).

#### Biotype and host plant

In field populations, *B. tabaci* biotypes were distributed differently among different host plants: the Q biotype was

dominant on cotton and sage in this study; and the B biotype was dominant on sweet pepper, cucumber, squash and eggplant (table 2). While such biotype-plant affinities may suggest specific adaptations, they can also reflect the insecticidal history of the crops. For instance, Israeli cotton fields are frequently treated with pesticides against B. tabaci during September, and this may favour the establishment of the insecticide-resistant Q biotype on that crop (Horowitz et al., 2005). Firm conclusions regarding the nature of possible biotype-plant associations await further investigation. Interestingly, all whiteflies sampled in the sage plots proved to be of the Q biotype and carried Rickettsia and Arsenophonus, a significantly higher infection rate than was found on other host plants. Although these SS may be advantageous for *B. tabaci* on this plant, a bias resulting from insecticide exposure cannot be ruled out. Controlled experiments with both biotypes on sage will be necessary to gain insight into the sage-Q biotype-Rickettsia and Arsenophonus tri-trophic interactions.

The importance of SS in the biology, ecology and evolution of *B. tabaci* biotypes will be clarified by further information regarding the phenotype of each SS.

#### Acknowledgements

The authors would like to thank Svetlana Kontsedalov for maintaining laboratory populations and Rami Horowitz for supplying whiteflies. Fabrice Vavré, Molly Hunter and two anonymous reviewers gave very constructive comments on early drafts of the manuscript. Thanks to Ruti Yonah for editing the manuscript. This research was supported by Grant No. 2004416 from the US-Israel Binational Science Foundation (BSF), Jerusalem, Israel to EZ-F and by the High Council for Scientific and Technological Cooperation between France and Israel to MG.

#### References

- Baumann, P. (2005) Biology of bacteriocyte-associated endosymbionts of plant sap-sucking insects. Annual Review of Microbiology 59, 155–189.
- Brown, J.K., Frohlich, D.R. & Rosell, R.C. (1995) The sweetpotato or silverleaf whiteflies – biotypes of *Bemisia tabaci* or a species complex. *Annual Review of Entomology* 40, 511– 534.
- Chen, D.Q., Montllor, C.B. & Purcell, A.H. (2000) Fitness effects of two facultative endosymbiotic bacteria on the pea aphid, *Acyrthosiphon pisum*, and the blue alfalfa aphid, *A. kondoi*. *Entomologia Experimentalis et Applicata* **95**, 315–323.
- Costa, H.S., Westcot, D.M., Ullman, D.E., Rosell, R.C., Brown, J.K. & Johnson, M.W. (1995) Morphological variation in *Bemisia* endosymbionts. *Protoplasma* 189, 194–202.
- Dale, C., Beeton, M., Harbison, C., Jones, T. & Pontes, M. (2006) Isolation, pure culture, and characterization of "Candidatus Arsenophonus arthropodicus," an intracellular secondary endosymbiont from the hippoboscid louse fly Pseudolynchia canariensis. Applied and Environmental Microbiology 72, 2997– 3004.
- Darby, A.C., Birkle, L.M., Turner, S.L. & Douglas, A.E. (2001) An aphid-borne bacterium allied to the secondary symbionts of whitefly. *FEMS Microbiology Ecology* 36, 43–50.
- Darby, A.C., Tosh, C.R., Walters, K.F.A. & Douglas, A.E. (2003) The significance of a facultative bacterium to natural

populations of the pea aphid *Acyrthosiphon pisum*. *Ecological Entomology* **28**, 145–150.

- De Barro, P.J. (2005) Genetic structure of the whitefly *Bemisia* tabaci in the Asia-Pacific region revealed using microsatellite markers. *Molecular Ecology* 14, 3695–3718.
- De Barro, P.J. & Hart, P.J. (2000) Mating interactions between two biotypes of the whitefly, *Bemisia tabaci* (Hemiptera: Aleyrodidae) in Australia. *Bulletin of Entomological Research* 90, 103–112.
- De Barro, P.J., Trueman, J.W.H. & Frohlich, D.R. (2005) Bemisia argentifolii is a race of B. tabaci (Hemiptera: Aleyrodidae): the molecular genetic differentiation of B. tabaci populations around the world. Bulletin of Entomological Research 95, 193–203.
- Everett, K.D.E., Thao, M.L., Horn, M., Dyszynski, G.E. & Baumann, P. (2005) Novel chlamydiae in whiteflies and scale insects: endosymbionts 'Candidatus Fritschea bemisiae' strain Falk and 'Candidatus Fritschea eriococci' strain Elm. International Journal of Systematic and Evolutionary Microbiology 55, 1581–1587.
- Ferrari, J., Darby, A.C., Daniell, T.J., Godfray, H.C.J. & Douglas, A.E. (2004) Linking the bacterial community in pea aphids with host-plant use and natural enemy resistance. *Ecological Entomology* 29, 60–65.
- Frohlich, D.R., Torres-Jerez, I., Bedford, I.D., Markham, P.G. & Brown, J.K. (1999) A phylogeographical analysis of the *Bemisia tabaci* species complex based on mitochondrial DNA markers. *Molecular Ecology* 8, 1683–1691.
- Gerling, D. & Mayer, R.T. (1996) Bemisia: 1995 Taxonomy, Biology, Damage, Control and Management. 702 pp. Andover, UK, Intercept Press.
- Gherna, R.L., Werren, J.H., Weisburg, W., Cote, R., Woese, C.R., Mandelco, L. & Brenner, D.J. (1991) Arsenophonusnasoniae gen-nov, sp-nov, the causative agent of the sonkiller trait in the parasitic wasp Nasonia vitripennis. International Journal of Systemic Bacteriology 41, 563–565.
- Gottlieb, Y., Ghanim, M., Chiel, E., Gerling, D., Portnoy, V., Steinberg, S., Tzuri, G., Horowitz, A.R., Belausov, E., Mozes-Daube, N., Kontsedalov, S., Gershon, M., Gal, S., Katzir, N. & Zchori-Fein, E. (2006) Identification and localization of a *Rickettsia* sp in *Bemisia tabaci* (Homoptera: Aleyrodidae). *Applied and Environmental Microbiology* 72, 3646–3652.
- Haynes, S., Darby, A.C., Daniell, T.J., Webster, G., Van Veen, F.J.F., Godfray, H.C.J., Prosser, J.I. & Douglas, A.E. (2003) Diversity of bacteria associated with natural aphid populations. *Applied and Environmental Microbiology* 69, 7216–7223.
- Heddi, A., Grenier, A.M., Khatchadourian, C., Charles, H. & Nardon, P. (1999) Four intracellular genomes direct weevil biology: Nuclear, mitochondrial, principal endosymbiont, and Wolbachia. Proceedings of the National Academy of Sciences USA 96, 6814–6819.
- Horowitz, A.R., Kontsedalov, S., Khasdan, V. & Ishaaya, I. (2005) Biotypes B and Q of *Bemisia tabaci* and their relevance to neonicotinoid and pyriproxyfen resistance. *Archives of Insect Biochemistry and Physiology* 58, 216–225.
- Khasdan, V., Levin, I., Rosner, A., Morin, S., Kontsedalov, S., Maslenin, L. & Horowitz, A.R. (2005) DNA markers for identifying biotypes B and Q of *Bemisia tabaci* (Hemiptera: Aleyrodidae) and studying population dynamics. *Bulletin of Entomological Research* 95, 605–613.
- Koga, R., Tsuchida, T. & Fukatsu, T. (2003) Changing partners in an obligate symbiosis: a facultative endosymbiont can compensate for loss of the essential endosymbiont *Buchmera*

in an aphid. Proceedings of the Royal Society Series B 270, 2543–2550.

- Leonardo, T.E. & Mondor, E.B. (2006) Symbiont modifies host life-history traits that affect gene flow. *Proceedings of the Royal Society Series B* 273, 1079–1084.
- Ma, D., Hadjistylli, M., Gorman, K., Denholm, I. & Devine, G. (2004) Pre- and post-zygotic breeding incompatibilities between B and Q biotypes of *Bemisia tabaci*. pp. 13–14 in *Proceedings of the 2nd European whitefly Symposium*. Cavtat, Croatia, 5–9 October 2004. (http://www.whitefly.org/ ews/ewsii-AbsCompendium.pdf)
- Montllor, C.B., Maxmen, A. & Purcell, A.H. (2002) Facultative bacterial endosymbionts benefit pea aphids Acyrthosiphon pisum under heat stress. Ecological Entomology 27, 189–195.
- Nirgianaki, A., Banks, G.K., Frohlich, D.R., Veneti, Z., Braig, H.R., Miller, T.A., Bedford, I.D., Markham, P.G., Savakis, C. & Bourtzis, K. (2003) Wolbachia infections of the whitefly Bemisia tabaci. Current Microbiology 47, 93–101.
- Nombela, G., Beitia, F. & Muniz, M. (2001) A differential interaction study of *Bemisia tabaci* Q-biotype on commercial tomato varieties with or without the *Mi* resistance gene, and comparative host responses with the B-biotype. *Entomologia Experimentalis et Applicata* **98**, 339–344.
- Oliveira, M.R.V., Henneberry, T.J. & Anderson, P. (2001) History, current status, and collaborative research projects for *Bemisia tabaci*. Crop Protection 20, 709–723.
- Oliver, K.M., Russell, J.A., Moran, N.A. & Hunter, M.S. (2003) Facultative bacterial symbionts in aphids confer resistance to parasitic wasps. *Proceedings of the National Academy of Sciences USA* 100, 1803–1807.
- Pascual, S. & Callejas, C. (2004) Intra- and interspecific competition between biotypes B and Q of *Bemisia tabaci* (Hemiptera: Aleyrodidae) from Spain. *Bulletin of Entomological Research* 94, 369–375.
- Perlman, S.J., Hunter, M.S. & Zchori-Fein, E. (2006) The emerging diversity of *Rickettsia*. Proceedings of the Royal Society Series B 273, 2097–2106 (DOI: 10.1098/rspb.2006. 3541).
- Perring, T.M. (2001) The Bemisia tabaci species complex. Crop Protection 20, 725–737.
- Sakurai, M., Koga, R., Tsuchida, T., Meng, X.Y. & Fukatsu, T. (2005) *Rickettsia* symbiont in the pea aphid *Acyrthosiphon pisum*: Novel cellular tropism, effect on host fitness, and interaction with the essential symbiont *Buchnera*. *Applied and Environmental Microbiology* **71**, 4069–4075.
- Sandstrom, J.P., Russell, J.A., White, J.P. & Moran, N.A. (2001) Independent origins and horizontal transfer of bacterial symbionts of aphids. *Molecular Ecology* 10, 217–228.

- Scarborough, C.L., Ferrari, J. & Godfray, H.C.J. (2005) Aphid protected from pathogen by endosymbiont. *Science* 310, 1781–1781.
- Simon, J.C., Carre, S., Boutin, M., Prunier-Leterme, N., Sabater-Munoz, B., Latorre, A. & Bournoville, R. (2003) Host-based divergence in populations of the pea aphid: insights from nuclear markers and the prevalence of facultative symbionts. *Proceedings of the Royal Society Series B* 270, 1703–1712.
- Stouthamer, R., Breeuwer, J.A.J. & Hurst, G.D.D. (1999) Wolbachia pipientis: Microbial manipulator of arthropod reproduction. Annual Review of Microbiology 53, 71–102.
- Thao, M.L.L. & Baumann, P. (2004) Evidence for multiple acquisition of Arsenophonus by whitefly species (Sternorrhyncha: Aleyrodidae). Current Microbiology 48, 140–144.
- Tsuchida, T., Koga, R., Shibao, H., Matsumoto, T. & Fukatsu, T. (2002) Diversity and geographic distribution of secondary endosymbiotic bacteria in natural populations of the pea aphid, Acyrthosiphon pisum. Molecular Ecology 11, 2123– 2135.
- Tsuchida, T., Koga, R. & Fukatsu, T. (2004) Host plant specialization governed by facultative symbiont. *Science* 303, 1989–1989.
- Van Lenteren, J.C. & Noldus, L.P.J.J. (1990) Whitefly-Plant Relationships: Behavioural and Ecological Aspects. pp. 47– 90 in Gerling, D. (Ed.) Whiteflies: their Bionomics, Pest Status and Management. Andover, UK, Intercept Press.
- Weeks, A.R., Velten, R. & Stouthamer, R. (2003) Incidence of a new sex-ratio-distorting endosymbiotic bacterium among arthropods. *Proceedings of the Royal Society Series B* 270, 1857–1865.
- Werren, J.H. & Windsor, D.M. (2000) Wolbachia infection frequencies in insects: evidence of a global equilibrium? Proceedings of the Royal Society Series B 267, 1277–1285.
- Wu, D., Daugherty, S.C., Van Aken, S.E., Pai, G.H., Watkins, K.L., Khouri, H., Tallon, L.J., Zaborsky, J.M., Dunbar, H.E., Tran, P.L., Moran, N.A. & Eisen, J.A. (2006) Metabolic complementarity and genomics of the dual bacterial symbiosis of sharpshooters. *PLoS Biology* 4, 1079– 1092.
- Zchori-Fein, E. & Brown, J.K. (2002) Diversity of prokaryotes associated with *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae). Annals of the Entomological Society of America 95, 711–718.

(Accepted 4 January 2007) © 2007 Cambridge University Press